

Discover ... The Collection

by Portsmouth museums



A male Black-faced Golden Bowerbird or Flame Bowerbird, native to the forests of New Guinea, Indonesia. It was prepared by James Gardner, who set up a taxidermy business in the early 1840's.

The company was based at Oxford Street, London. Gardner held taxidermy warrants for the royal families of both England and France, as well as the Khedive of Egypt.

Discover ... The Collection

by Portsmouth museums



The Margay is adapted to life in the trees by being able to rotate its hind limbs 180 degrees. They are found in Mexico and Central and South America, but due to habitat loss, illegal hunting and trade, their populations have decreased.

Discover ... The Collection

by Portsmouth museums



Stag beetles are an impressive species which can be found mainly in the south east of the UK, and are often found in Hampshire.

They are the largest beetle species in the UK.

They get their name from the mandibles (jaws) of the males which look like two large antlers. These mandibles are used to fend off rival males.

Discover ... The Collection

by Portsmouth museums



Purple Herons are slightly smaller than the more familiar grey heron which you can see opposite. They are found on mainland Europe, but occasionally seen in the UK.

Discover ... The Collection

by Portsmouth museums



Badgers are omnivorous mammals found throughout the UK. They forage for earthworms, other invertebrates and fruit in the cover of darkness using their keen sense of smell and bulky bodies to dig out their food.